

**CH302H – Principles of Chemistry II: Honors**  
Fall 2016, Unique 49420

**Homework, Week 2**

1. Calculate the molar entropy of a constant-volume sample of neon at 500 K given that it is  $146.22 \text{ J K}^{-1} \text{ mol}^{-1}$  at 298 K and  $C_{V,m} = 3/2R$ .
2. Determine  $\Delta S$  (for the system) when 3.0 mol of an ideal gas at  $25^\circ\text{C}$  and 1.0 atm is heated to  $125^\circ\text{C}$  and expanded to 5.0 atm. Rationalize the sign of  $\Delta S$ .  $C_{V,m} = 3/2R$ ,  $C_{P,m} = 5/2R$ .
3. Consider a system containing 2.0 mol  $\text{CO}_2(\text{g})$  ( $C_{v,m} = 28.8 \text{ J K}^{-1} \text{ mol}^{-1}$ ), initially at  $25^\circ\text{C}$  and 10 atm and confined to a cylinder of cross-section  $10.0 \text{ cm}^2$ . It is allowed to expand adiabatically against a constant external pressure of 1.0 atm until the piston has moved outwards through 20 cm. Determine  $q$ ,  $w$ ,  $\Delta U$ ,  $\Delta H$ , and  $\Delta S$ .
4. Determine the standard reaction entropy at 298 K of the following reactions:
  - a)  $2 \text{ CH}_3\text{CHO}(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2 \text{ CH}_3\text{COOH}(\text{l})$
  - b)  $\text{Hg}(\text{l}) + \text{Cl}_2(\text{g}) \rightarrow \text{HgCl}_2(\text{s})$
5. A 500 g block of copper ( $C_{p,m} = 24.4 \text{ J K}^{-1} \text{ mol}^{-1}$ ) initially at 293 K is in thermal equilibrium with an electric heater of resistance 1.0 k $\Omega$  and negligible mass. A current of 1.0 A is passed for 15.0 s. Determine the change in entropy of the copper block. The experiment is then repeated with the copper immersed in a stream of water that maintains its temperature at 293 K. Determine the change in entropy of the copper and the water.
6. 4.0 moles of an ideal gas is divided evenly in a cylinder separated into 2 chambers, A and B, that are partitioned with a movable insulating boundary. Initially, the volume and temperature of both chambers is 2.0 L and 300 K, respectively. Each chamber has an independent heater that is used to supply heat. The heater for chamber B is used to maintain the temperature of B at all times. The heater for chamber A is used to supply heat to chamber A to move the boundary between A and B reversibly to decrease the volume of chamber B by half. Determine  $\Delta H$  for each chamber, assuming  $C_{V,m} = 20 \text{ J K}^{-1} \text{ mol}^{-1}$ .
7. For each of the following processes determine whether  $\Delta S_{\text{sys}}$  is greater than zero, less than zero, or equal to zero. Explain your reasoning.
  - a) A process in which no heat is exchanged between system and surroundings
  - b) An isothermal expansion of an ideal gas
  - c) An isobaric (i.e constant pressure) cooling of an ideal gas
  - d) Isobaric evaporation of a liquid.

8. Vaporization at the normal boiling point of a substance (the boiling point at 1 atm,  $T_{vap}$ ) is a reversible process. If  $\Delta H_{vap}$  of water is  $40.65 \text{ kJ mol}^{-1}$ , determine  $\Delta S_{vap}$  when 2.0 moles of water are vaporized at  $100^\circ\text{C}$ . Comment on the sign of  $\Delta S_{vap}$ .